

# Foundation and generalization of the expansion by regions

**Bernd Jantzen**

*RWTH Aachen University*


arXiv:1111.2589 [hep-ph]

- I The strategy of regions
- II Why does the method work?
- III Examples
- IV The general formalism
- V Summary

# I The strategy of regions

## Starting point: (multi-)loop integral

$$F = \int d^d k_1 \int d^d k_2 \cdots \frac{1}{(k_1 + p_1)^2 - m_1^2} \times \\ \times \frac{1}{(k_1 + k_2 + p_2)^2 - m_2^2} \cdots$$



- complicated function of internal masses  $m_i$  and kinematical parameters  $p_i^2$ ,  $p_i \cdot p_j$
- exact evaluation often hard or impossible

**Exploit parameter hierarchies**, e.g. large energies  $Q \gg$  small masses  $m$ :

↪ **expand integral** in small ratios  $\frac{m^2}{Q^2}$

↪ simplification achieved if **expansion of integrand before integration**


**But:**

- ★ loop-momentum components  $k_i^\mu$  can take any values (large, small, mixed, ...)
- ★ naive expansions of integrand may **generate new singularities**
- ↪ Need sophisticated methods of **asymptotic expansions**.

## Simple example: large-momentum expansion

$$F = \int \frac{Dk}{(k+p)^2 (k^2 - m^2)^2} \quad \left[ \int Dk \equiv \mu^{2\epsilon} e^{\epsilon\gamma_E} \int \frac{d^d k}{i\pi^{d/2}} \right]$$

$$d = 4 - 2\epsilon$$



Large momentum  $|p^2| \gg m^2 \rightsquigarrow$  expand in  $\frac{m^2}{p^2}$ .

Integral is UV- and IR-finite, the exact result is known:

$$[p^2 \rightarrow p^2 + i0]$$

$$F = \frac{1}{p^2} \left[ \ln\left(\frac{-p^2}{m^2}\right) + \ln\left(1 - \frac{m^2}{p^2}\right) \right] + \mathcal{O}(\epsilon)$$

$$\xrightarrow{\text{expand}} \frac{1}{p^2} \left[ \ln\left(\frac{-p^2}{m^2}\right) - \sum_{j=1}^{\infty} \frac{1}{j} \left(\frac{m^2}{p^2}\right)^j \right] + \mathcal{O}(\epsilon)$$


Now assume that we could not calculate this integral exactly ...

## Large-momentum expansion (2)

Large momentum  $|p^2| \gg m^2$

↪ expand integrand before integration:

$$F = \int \frac{Dk}{(k+p)^2 (k^2 - m^2)^2}$$



## Expansion by regions

↪ here 2 relevant **regions**:

**Beneke, Smirnov, Nucl. Phys. B 522, 321 (1998)**

**Smirnov, Rakhmetov, Theor. Math. Phys. 120, 870 (1999)**

**Smirnov, Phys. Lett. B 465, 226 (1999)**

- **hard (h):**  $k \sim p \Rightarrow \sum_i T_i^{(h)} \frac{1}{(k^2 - m^2)^2} = \sum_{i=0}^{\infty} (1+i) \frac{(m^2)^i}{(k^2)^{2+i}}$

- **soft (s):**  $k \sim m \Rightarrow \sum_j T_j^{(s)} \frac{1}{(k+p)^2} = \sum_{j_1, j_2=0}^{\infty} \frac{(j_1 + j_2)!}{j_1! j_2!} \frac{(-2k \cdot p)^{j_1} (-k^2)^{j_2}}{(p^2)^{1+j_1+j_2}}$

⇒ Integrate each expanded term over the **whole integration domain**.

⇒ Set scaleless integrals to zero (like in dimensional regularization).

## Leading-order contributions:

- **hard:**  $F_0^{(h)} = \int \frac{Dk}{(k+p)^2 (k^2)^2} = \frac{1}{p^2} \left( \frac{\mu^2}{-p^2} \right)^\epsilon \left( -\frac{1}{\epsilon} + \mathcal{O}(\epsilon) \right)$

- **soft:**  $F_0^{(s)} = \int \frac{Dk}{p^2 (k^2 - m^2)^2} = \frac{1}{p^2} \left( \frac{\mu^2}{m^2} \right)^\epsilon \left( \frac{1}{\epsilon} + \mathcal{O}(\epsilon) \right)$


↪ Contributions are **homogeneous** functions of the expansion parameter  $\frac{m^2}{p^2}$ .

## Large-momentum expansion (3)

### Leading-order contributions:

- **hard:**  $F_0^{(h)} = \frac{1}{p^2} \left[ -\frac{1}{\epsilon} + \ln\left(\frac{-p^2}{\mu^2}\right) \right] + \mathcal{O}(\epsilon) \rightsquigarrow \text{IR-singular!}$

- **soft:**  $F_0^{(s)} = \frac{1}{p^2} \left[ \frac{1}{\epsilon} + \ln\left(\frac{\mu^2}{m^2}\right) \right] + \mathcal{O}(\epsilon) \rightsquigarrow \text{UV-singular!}$



↪ Singularities are cancelled in the sum of all contributions, exact result approximated:

$$F_0 = F_0^{(h)} + F_0^{(s)} = \frac{1}{p^2} \ln\left(\frac{-p^2}{m^2}\right) + \mathcal{O}(\epsilon) = F + \mathcal{O}\left(\frac{m^2}{(p^2)^2}\right) \quad \checkmark$$

### Expand to all orders in $\frac{m^2}{p^2}$ :

$$[(\alpha)_n = \Gamma(\alpha + n)/\Gamma(\alpha)]$$

$$F^{(h)} = \frac{1}{p^2} \left(\frac{\mu^2}{-p^2}\right)^\epsilon \frac{e^{\epsilon\gamma_E} \Gamma(1+\epsilon) \Gamma^2(1-\epsilon)}{(-\epsilon) \Gamma(1-2\epsilon)} \sum_{i=0}^{\infty} \left(\frac{m^2}{p^2}\right)^i \frac{(2\epsilon)_i}{i!} = F_0^{(h)} + \frac{2}{p^2} \ln\left(1 - \frac{m^2}{p^2}\right) + \mathcal{O}(\epsilon)$$

$$F^{(s)} = \frac{1}{p^2} \left(\frac{\mu^2}{m^2}\right)^\epsilon e^{\epsilon\gamma_E} \Gamma(\epsilon) \sum_{j=0}^{\infty} \left(\frac{m^2}{p^2}\right)^j \frac{(\epsilon)_j}{(1-\epsilon)_j} = F_0^{(s)} - \frac{1}{p^2} \ln\left(1 - \frac{m^2}{p^2}\right) + \mathcal{O}(\epsilon)$$

$$\hookrightarrow F = F^{(h)} + F^{(s)} = \frac{1}{p^2} \left[ \ln\left(\frac{-p^2}{m^2}\right) + \ln\left(1 - \frac{m^2}{p^2}\right) \right] + \mathcal{O}(\epsilon) \quad \checkmark$$

⇒ Full result  $F$  exactly reproduced.

## Questions: Why does this expansion by regions work?


- Didn't we **double-count** every  $k \in \mathbb{R}^d$  when replacing  $\int \mathcal{D}k \rightarrow \int \mathcal{D}k T_0^{(h)} + \int \mathcal{D}k T_0^{(s)}$  ?
- What ensures the **cancellation of singularities**? (IR  $\leftrightarrow$  UV!)
- How do we know that the chosen **set of regions** is **complete**?
- What is the role of **scaleless integrals**?

## II Why does the method work?

Idea based on a 1-dimensional toy example from M. Beneke (1997)  
[see also: Smirnov, *Applied Asymptotic Expansions In Momenta And Masses*]

### Large-momentum example

Let us show step by step how the expansions reproduce the full result.



The **expansions**  $\sum_i T_i^{(h)}$ ,  $\sum_j T_j^{(s)}$  **converge absolutely** within **domains**  $D_h$ ,  $D_s$ :

$$(h): \frac{1}{(k^2 - m^2)^2} = \sum_i T_i^{(h)} \frac{1}{(k^2 - m^2)^2} \text{ within } D_h = \left\{ k \in \mathbb{R}^d : |k^2| \geq \Lambda^2 \right\},$$

$$(s): \frac{1}{(k+p)^2} = \sum_j T_j^{(s)} \frac{1}{(k+p)^2} \text{ within } D_s = \left\{ k \in \mathbb{R}^d : |k^2| < \Lambda^2 \right\},$$


with  $m^2 \ll \Lambda^2 \ll |p^2| \rightsquigarrow D_h \cup D_s = \mathbb{R}^d$ ,  $D_h \cap D_s = \emptyset$ .

The expansions **commute** with **integrals restricted to the corresponding domains**:

$$F = \int_{k \in \mathbb{R}^d} \underbrace{\frac{1}{(k+p)^2 (k^2 - m^2)^2}}_I = \sum_i \int_{k \in D_h} \text{D}k T_i^{(h)} I + \sum_j \int_{k \in D_s} \text{D}k T_j^{(s)} I$$

Continue transforming the expression for the full integral:

$$\begin{aligned}
 F &= \int_{k \in \mathbb{R}^d} \mathrm{D}k \underbrace{\frac{1}{(k+p)^2 (k^2 - m^2)^2}}_I = \sum_i \int_{k \in D_h} \mathrm{D}k T_i^{(h)} I + \sum_j \int_{k \in D_s} \mathrm{D}k T_j^{(s)} I \\
 &= \sum_i \left( \int_{k \in \mathbb{R}^d} \mathrm{D}k T_i^{(h)} I - \sum_j \int_{k \in D_s} \mathrm{D}k T_j^{(s)} T_i^{(h)} I \right) + \sum_j \left( \int_{k \in \mathbb{R}^d} \mathrm{D}k T_j^{(s)} I - \sum_i \int_{k \in D_h} \mathrm{D}k T_i^{(h)} T_j^{(s)} I \right)
 \end{aligned}$$




The **expansions commute**:  $T_i^{(h)} T_j^{(s)} I = T_j^{(s)} T_i^{(h)} I \equiv T_{i,j}^{(h,s)} I$

$$\Rightarrow \text{Identity: } F = \underbrace{\sum_i \int \mathrm{D}k T_i^{(h)} I}_{F^{(h)}} + \underbrace{\sum_j \int \mathrm{D}k T_j^{(s)} I}_{F^{(s)}} - \underbrace{\sum_{i,j} \int \mathrm{D}k T_{i,j}^{(h,s)} I}_{F^{(h,s)}}$$

All terms integrated over the **whole integration domain**  $\mathbb{R}^d$  as prescribed for the expansion by regions  $\Rightarrow$  location of **boundary**  $\Lambda$  between  $D_h, D_s$  **irrelevant**.



**Identity:** 
$$F = \underbrace{\sum_i \int Dk T_i^{(h)} I}_{F^{(h)}} + \underbrace{\sum_j \int Dk T_j^{(s)} I}_{F^{(s)}} - \underbrace{\sum_{i,j} \int Dk T_{i,j}^{(h,s)} I}_{F^{(h,s)}}$$



**Additional overlap contribution  $F^{(h,s)}$ ?**

$$F^{(h,s)} = \sum_{i=0}^{\infty} (1+i) \sum_{j_1, j_2=0}^{\infty} (-1)^{j_2} \frac{(j_1 + j_2)!}{j_1! j_2!} \frac{(m^2)^i}{(p^2)^{1+j_1+j_2}} \int Dk \frac{(-2k \cdot p)^{j_1}}{(k^2)^{2+i-j_2}} = 0 \quad \text{scaleless!}$$

[Actually  $\int \frac{Dk}{(k^2)^2} = \frac{1}{\epsilon_{UV}} - \frac{1}{\epsilon_{IR}}$  cancels corresponding singularities in  $F^{(h)}$  and  $F^{(s)}$ .]

$\hookrightarrow \boxed{F = F^{(h)} + F^{(s)}}$  as found before.

But now this identity has been obtained **without evaluating  $F, F^{(h)}, F^{(s)}$ !**

### III Examples


#### Example with 3 regions: threshold expansion for heavy-particle pair production

Regions analyzed in Beneke, Smirnov, NPB 522, 321 (1998)

Centre-of-mass system:  $(q^\mu) = (q_0, \vec{0})$ ,  $(p^\mu) = (0, \vec{p})$

Close to threshold:  $q^2 \approx (2m)^2 \Rightarrow q^2 \gg |p^2|$  or  $q_0 \gg |\vec{p}|$

$$F = \int \frac{Dk}{(k^2 + q_0 k_0 - 2\vec{p} \cdot \vec{k})(k^2 - q_0 k_0 - 2\vec{p} \cdot \vec{k}) k^2}$$



Relevant regions:


- **hard** ( $h$ ):  $k_0, |\vec{k}| \sim q_0 \Rightarrow$  expansion  $\sum_j T_j^{(h)}$  converges in  $D_h$
- **soft** ( $s$ ):  $k_0, |\vec{k}| \sim |\vec{p}| \Rightarrow$  expansion  $\sum_j T_j^{(s)}$  converges in  $D_s$
- **potential** ( $p$ ):  $k_0 \sim \frac{\vec{p}^2}{q_0}$ ,  $|\vec{k}| \sim |\vec{p}| \Rightarrow$  expansion  $\sum_j T_j^{(p)}$  converges in  $D_p$

$$\hookrightarrow D_h \cup D_s \cup D_p = \mathbb{R}^d, \quad D_h \cap D_s = D_h \cap D_p = D_s \cap D_p = \emptyset$$

$\hookrightarrow$  The expansions  $T^{(h)}, T^{(s)}, T^{(p)}$  commute with each other.

## Threshold expansion (2)

Similar transformations as for the large-momentum example yield the following **identity**:



$$F = F^{(h)} + \underbrace{F^{(s)}}_{=0} + F^{(p)} - \left( \underbrace{F^{(h,s)}}_{=0} + \underbrace{F^{(h,p)}}_{=0} + \underbrace{F^{(s,p)}}_{=0} \right) + \underbrace{F^{(h,s,p)}}_{=0 \text{ (scaleless)}}$$

with

$$F^{(h)} = -\frac{2}{q^2} \left( \frac{4\mu^2}{q^2} \right)^\epsilon e^{\epsilon\gamma_E} \Gamma(\epsilon) \sum_{j=0}^{\infty} \left( -\frac{4p^2}{q^2} \right)^j \frac{(1+\epsilon)_j}{j! (1+2\epsilon+2j)}$$

$$F^{(p)} = \frac{e^{\epsilon\gamma_E} \Gamma(\frac{1}{2} + \epsilon) \sqrt{\pi}}{2\epsilon \sqrt{q^2 (p^2 - i0)}} \left( \frac{\mu^2}{p^2 - i0} \right)^\epsilon \quad [\text{higher orders are scaleless}]$$


Exact result reproduced:

$$F^{(h)} + F^{(p)} = F = \frac{e^{\epsilon\gamma_E} \Gamma(\epsilon)}{2p^2} \left( \frac{\mu^2}{p^2 - i0} \right)^\epsilon {}_2F_1 \left( \frac{1}{2}, 1 + \epsilon; \frac{3}{2}; -\frac{q^2}{4p^2} - i0 \right) \quad \checkmark$$

## Example with non-commuting expansions: Sudakov form factor

Cannot always choose expansions which commute with each other.

Sudakov limit:  $-(p_1 - p_2)^2 = \boxed{Q^2 \gg m^2}$




$$F = \int \frac{Dk}{(k^+k^- - \vec{k}_\perp^2 + Qk^+)^{1+\delta} (k^+k^- - \vec{k}_\perp^2 + Qk^-)^{1-\delta} (k^+k^- - \vec{k}_\perp^2 - m^2)}$$

$\hookrightarrow$  analytic regulator  $\delta \rightarrow 0$

[light-cone coordinates:  $2p_{1,2} \cdot k = Qk^\pm$ ,  $p_{1,2} \cdot k_\perp = 0$ ]

### Regions & domains:

- **hard (h)**:  $k^+, k^-, |\vec{k}_\perp| \sim Q \Rightarrow D_h = \{k \in \mathbb{R}^d : \vec{k}_\perp^2 \gg m^2\}$
- **1-collinear (1c)**:  $k^+ \sim \frac{m^2}{Q}$ ,  $k^- \sim Q$ ,  $|\vec{k}_\perp| \sim m$
- **2-collinear (2c)**:  $k^+ \sim Q$ ,  $k^- \sim \frac{m^2}{Q}$ ,  $|\vec{k}_\perp| \sim m$
- **Glauber (g)**:  $k^+, k^- \sim \frac{m^2}{Q}$ ,  $|\vec{k}_\perp| \sim m$
- **collinear-plane (cp)**:  $k^+, k^- \sim Q$ ,  $|\vec{k}_\perp| \sim m$   
 $\hookrightarrow$  “artificial” region to ensure  $\cup_x D_x = \mathbb{R}^d$



[No soft region needed:  $T^{(s)} \equiv T^{(1c)}T^{(2c)}$ ]

Most expansions commute, but  $T^{(g)}T^{(cp)} \neq T^{(cp)}T^{(g)}$  !

## Sudakov form factor (2)

$T^{(g)}T^{(cp)} \neq T^{(cp)}T^{(g)} \rightsquigarrow$  Construct **identity** avoiding combination of  $(g)$  and  $(cp)$ :

$$\begin{aligned}
 F &= F^{(h)} + F^{(1c)} + F^{(2c)} + F^{(g)} + F^{(cp)} \\
 &- \left( F^{(h,1c)} + F^{(h,2c)} + F^{(h,g)} + F^{(h,cp)} + F^{(1c,2c)} + F^{(1c,g)} + F^{(1c,cp)} + F^{(2c,g)} + F^{(2c,cp)} \right) \\
 &+ F^{(h,1c,2c)} + F^{(h,1c,g)} + F^{(h,1c,cp)} + F^{(h,2c,g)} + F^{(h,2c,cp)} + F^{(1c,2c,g)} + F^{(1c,2c,cp)} \\
 &- \left( F^{(h,1c,2c,g)} + F^{(h,1c,2c,cp)} \right) + F_{cp \leftarrow g}^{\text{extra}} + F_{g \leftarrow cp}^{\text{extra}}
 \end{aligned}$$

### Usual terms:

- no combination of  $(g)$  and  $(cp)$
- $F^{(g)}$ ,  $F^{(cp)}$  and all overlap contributions are scaleless (with analytic regularization)


### Extra terms:

- $F_{cp \leftarrow g}^{\text{extra}}$  involves  $T^{(cp)}T^{(g)}$  integrated over  $k \in D_{cp}$
- $F_{g \leftarrow cp}^{\text{extra}}$  involves  $T^{(g)}T^{(cp)}$  integrated over  $k \in D_g$

Both extra terms cancel at the integrand level,

e.g. in  $F_{g \leftarrow cp}^{\text{extra}}$  because  $T^{(x)}T^{(g)}T^{(cp)} = T^{(g)}T^{(cp)} \forall x \in \{h, 1c, 2c\}$ .


[They must cancel  $\rightsquigarrow$  otherwise dependence on boundaries of  $D_g$ ,  $D_{cp}$ .]



## Sudakov form factor (3)

Omitting scaleless contributions and vanishing extra terms:

$$F = F^{(h)} + F^{(1c)} + F^{(2c)}$$



Regions explicitly evaluated to all orders in  $\frac{m^2}{Q^2}$ :

[omitting  $\mathcal{O}(\delta)$  and  $\mathcal{O}(\epsilon)$ ]

$$F^{(h)} = -\frac{1}{Q^2} \left( \frac{\mu^2}{Q^2} \right)^\epsilon \left\{ \frac{1}{\epsilon^2} - \frac{2}{\epsilon} \ln \left( 1 - \frac{m^2}{Q^2} \right) + \ln^2 \left( 1 - \frac{m^2}{Q^2} \right) - 2 \text{Li}_2 \left( \frac{m^2}{Q^2} \right) - \frac{\pi^2}{12} \right\}$$

$$F^{(1c)}, F^{(2c)} = -\frac{1}{2Q^2} \left( \frac{\mu^2}{Q^2} \right)^\epsilon \left\{ \pm \frac{1}{\delta} \left[ \frac{1}{\epsilon} + \ln \frac{Q^2}{m^2} - \ln \left( 1 - \frac{m^2}{Q^2} \right) \right] - \frac{1}{\epsilon^2} + \frac{2}{\epsilon} \ln \left( 1 - \frac{m^2}{Q^2} \right) + \frac{1}{2} \ln^2 \frac{Q^2}{m^2} + \ln \frac{Q^2}{m^2} \ln \left( 1 - \frac{m^2}{Q^2} \right) - \ln^2 \left( 1 - \frac{m^2}{Q^2} \right) + \text{Li}_2 \left( \frac{m^2}{Q^2} \right) + \frac{5}{12} \pi^2 \right\}$$

$\hookrightarrow F^{(1c)}$  and  $F^{(2c)}$  are **not separately finite for  $\delta \rightarrow 0$** , but their sum is.

Compare to exact result:

$$F = -\frac{1}{Q^2} \left\{ \frac{1}{2} \ln^2 \frac{Q^2}{m^2} + \ln \frac{Q^2}{m^2} \ln \left( 1 - \frac{m^2}{Q^2} \right) - \text{Li}_2 \left( \frac{m^2}{Q^2} \right) + \frac{\pi^2}{3} \right\} \quad \checkmark$$

## IV The general formalism

Identities as in the previous examples are **generally valid**, under some conditions.

### Consider

- a (multiple) integral  $F = \int Dk I$  over the domain  $D$  (e.g.  $D = \mathbb{R}^d$ ),
- a set of  $N$  regions  $R = \{x_1, \dots, x_N\}$ ,
- for each region  $x \in R$  an expansion  $T^{(x)} = \sum_j T_j^{(x)}$  which converges absolutely in the domain  $D_x \subset D$ .

### Conditions

- $\bigcup_{x \in R} D_x = D$ ,  $D_x \cap D_{x'} = \emptyset \forall x \neq x'$ .
- Some of the **expansions commute** with each other.  
Let  $R_c = \{x_1, \dots, x_{N_c}\}$  and  $R_{nc} = \{x_{N_c+1}, \dots, x_N\}$  with  $1 \leq N_c \leq N$ .  
Then:  $T^{(x)} T^{(x')} = T^{(x')} T^{(x)} \equiv T^{(x, x')} \forall x \in R_c, x' \in R$ .
- Every pair of non-commuting expansions is invariant under some expansion from  $R_c$ :  
 $\forall x'_1, x'_2 \in R_{nc} \exists x \in R_c : T^{(x)} T^{(x'_2)} T^{(x'_1)} = T^{(x'_2)} T^{(x'_1)}$ .
- $\exists$  **regularization** for singularities, e.g. dimensional (+ analytic) regularization.  
 $\hookrightarrow$  All expanded integrals and series expansions in the formalism are well-defined.

## The general formalism (2)

Under these conditions, the following **identity** holds:  $[F^{(x,\dots)} \equiv \sum_{j,\dots} \int Dk T_{j,\dots}^{(x,\dots)} I]$

$$F = \sum_{x \in R} F^{(x)} - \sum_{\{x'_1, x'_2\} \subset R}^{\langle R_c + 1 \rangle} F^{(x'_1, x'_2)} + \dots - (-1)^n \sum_{\{x'_1, \dots, x'_n\} \subset R}^{\langle R_c + 1 \rangle} F^{(x'_1, \dots, x'_n)} + \dots + (-1)^{N_c} \sum_{x' \in R_{nc}} F^{(x', x_1, \dots, x_{N_c})}$$


where the sums run over subsets  $\{x'_1, \dots\}$  containing at most one region from  $R_{nc}$ .

## Comments

- This identity is **exact** when the expansions are summed to all orders. ✓  
Leading-order approximation for  $F \rightsquigarrow$  dropping higher-order terms.
- It is **independent of the regularization** (dim. reg., analytic reg., cut-off, infinitesimal masses/off-shellness, ...) as long as all individual terms are well-defined.
- Usually regions & regularization are chosen such that **multiple expansions**  $F^{(x'_1, \dots, x'_n)}$  ( $n \geq 2$ ) are **scaleless** and vanish.  
[✓ if each  $F_0^{(x)}$  is a *homogeneous* function of the expansion parameter with *unique scaling*.]
- If  $\exists F^{(x'_1, x'_2, \dots)} \neq 0 \rightsquigarrow$  relevant **overlap contributions** ( $\rightarrow$  “**zero-bin subtractions**”).  
They appear e.g. when avoiding analytic regularization in SCET. e.g. Manohar, Stewart '06;  
Chiu, Fuhrer, Hoang, Kelley, Manohar '09; ...



## Example with relevant overlap contributions: forward scattering with small momentum exchange



Two light-like particles with large center-of-mass energy exchange a small momentum  $r$ :



$$p_1^2 = (p_1 - r)^2 = p_2^2 = (p_2 + r)^2 = 0$$

$$(p_1 + p_2)^2 = \boxed{Q^2 \gg \vec{r}_\perp^2}, \quad r^\pm \approx \mp \frac{\vec{r}_\perp^2}{Q}$$

Symmetrize integral under  $k \leftrightarrow r - k$

$\hookrightarrow$  avoids divergences at  $|k^\pm| \rightarrow \infty$  under expansion.

$$F = \frac{1}{2} \int \frac{Dk}{k^2 (r - k)^2} \left( \frac{1}{((p_1 - k)^2)^{1+\delta}} + \frac{1}{((p_1 - r + k)^2)^{1+\delta}} \right) \times \left( \frac{1}{((p_2 + k)^2)^{1-\delta}} + \frac{1}{((p_2 + r - k)^2)^{1-\delta}} \right)$$




**Regions:** same as for Sudakov form factor (scaling with  $m \rightarrow |\vec{r}_\perp|$ ),

**Domains:** similar (but more involved for  $|\vec{k}_\perp| \gg |\vec{r}_\perp|$ )

## Forward scattering (2)

Same identity as for Sudakov form factor:



$$\begin{aligned}
 F &= F^{(h)} + F^{(1c)} + F^{(2c)} + F^{(g)} + F^{(cp)} \\
 &\quad - \left( F^{(h,1c)} + F^{(h,2c)} + F^{(h,g)} + F^{(h,cp)} + F^{(1c,2c)} + F^{(1c,g)} + F^{(1c,cp)} + F^{(2c,g)} + F^{(2c,cp)} \right) \\
 &\quad + F^{(h,1c,2c)} + F^{(h,1c,g)} + F^{(h,1c,cp)} + F^{(h,2c,g)} + F^{(h,2c,cp)} + F^{(1c,2c,g)} + F^{(1c,2c,cp)} \\
 &\quad - \left( F^{(h,1c,2c,g)} + F^{(h,1c,2c,cp)} \right)
 \end{aligned}$$

**With analytic regulator  $\delta \rightarrow 0$ :**  $F_0 = F_0^{(1c)} + F_0^{(2c)}$  [ $F_0^{(h)}$  suppressed, others scaleless]

$$F_0^{(1c)} = F_0^{(2c)} = \frac{1}{2} \frac{i\pi}{Q^2 \vec{r}_\perp^2} \left( \frac{\mu^2}{\vec{r}_\perp^2} \right)^\epsilon \frac{e^{\epsilon\gamma_E} \Gamma(1+\epsilon) \Gamma^2(-\epsilon)}{\Gamma(-2\epsilon)}$$

**Without analytic regularization ( $\delta = 0$ ):** [all terms are still well-defined]

$$F_0 = F_0^{(1c)} + F_0^{(2c)} + F_0^{(g)} - \left( F_0^{(1c,2c)} + F_0^{(1c,g)} + F_0^{(2c,g)} \right) + F_0^{(1c,2c,g)}$$

$$F_0^{(x,\dots)} = \frac{i\pi}{Q^2 \vec{r}_\perp^2} \left( \frac{\mu^2}{\vec{r}_\perp^2} \right)^\epsilon \frac{e^{\epsilon\gamma_E} \Gamma(1+\epsilon) \Gamma^2(-\epsilon)}{\Gamma(-2\epsilon)} \quad \forall x, \dots \in \{1c, 2c, g\}$$

$\hookrightarrow$  consistent results independent of regularization:  $\frac{1}{2} + \frac{1}{2} = 1 + 1 + 1 - (1 + 1 + 1) + 1 \checkmark$

$\hookrightarrow$  agreement with leading-order expansion of full result

# V Summary

## Expansion by regions for general integrals

- **Conditions for regions** (+ corresponding expansions & domains) established.
- **Identity proven**  $\rightsquigarrow$  relates exact integral to sum of expanded terms:

$$F = \sum_{x \in R} F^{(x)} - \sum_{\{x'_1, x'_2\} \subset R}^{\langle R_c + 1 \rangle} F^{(x'_1, x'_2)} + \dots - (-1)^n \sum_{\{x'_1, \dots, x'_n\} \subset R}^{\langle R_c + 1 \rangle} F^{(x'_1, \dots, x'_n)} + \dots + (-1)^{N_c} \sum_{x' \in R_{nc}} F^{(x', x_1, \dots, x_{N_c})}$$

- $\hookrightarrow$  valid **independent of the choice of regularization**
- This identity includes **overlap contributions** with multiple expansions
  - $\hookrightarrow$  can be **scaleless**  $\rightsquigarrow$  known recipe for expansion by regions ✓  
or **relevant** (depending on regularization)
  - $\hookrightarrow$  **generalization** of known recipe.

## Application to example integrals

- setup of the regions, expansions & convergence domains,
- check of conditions,
- evaluation of expanded integrals,
- comparison to exact result.

**Extra slides**

## “Real-life” example

The expansion by regions has been applied successfully to many complicated loop integrals.

Example:


Denner, B.J., Pozzorini '08

### 2-loop vertex integral in the high-energy limit

$Q^2 \gg m_t^2 \sim M_{W,Z}^2 \Rightarrow 9$  relevant regions: [labelled “ $(k_1 - k_2)$ ”]

$(h - h), (1c - h), (h - 2c), (1c - 1c), (1c - 2c),$   
 $(2c - 2c), (us - 2c), (1c - 2uc), (2uc - 2uc)$

$\hookrightarrow$  Next-to-leading-logarithmic result obtained and cross-checked with other methods.



## Practical note: how to find the relevant regions

- Look where the **propagators** have **poles**:
  - ★ Large-momentum example:  $(k + p)^2 = 0$  at  $k \sim p$ ,  $k^2 - m^2 = 0$  at  $k \sim m$ .
  - ★ Close the integration contour of one component (e.g.  $k^0$ ,  $k^\pm$ ).  
For all residues investigate the scaling of the components.
- Use **Mellin–Barnes (MB) representations**:
  1. Evaluate the full (scalar) integral for general propagator powers  $n_i$  in terms of multiple MB integrals.
  2. Close MB contours involving the expansion parameter and extract the leading contributions.
  3. The individual terms can be identified with corresponding regions by their homogeneous scaling with the expansion parameter depending on  $d$  and  $n_i$ .  
[A subsequent expansion by regions often yields simpler expressions for the contributions.]
- **Try all possible regions** that you can imagine ...  
If a region does not contribute, its integrals are scaleless.
- When a region is missing, the total result is often (but not always) more singular than it should be.  $\rightsquigarrow$  Important cross-check, but no guarantee!

## Threshold expansion for heavy-particle pair production


[details of  $D_h, D_s, D_p$ ]

Regions analyzed in Beneke, Smirnov, NPB 522, 321 (1998)

Centre-of-mass system:  $(q^\mu) = (q_0, \vec{0})$ ,  $(p^\mu) = (0, \vec{p})$

Close to threshold:  $q^2 \approx (2m)^2 \Rightarrow q^2 \gg |p^2|$  or  $q_0 \gg |\vec{p}|$

$$F = \int \frac{Dk}{(k^2 + q_0 k_0 - 2\vec{p} \cdot \vec{k})(k^2 - q_0 k_0 - 2\vec{p} \cdot \vec{k}) k^2}$$



Relevant regions:

- **hard** ( $h$ ):  $k_0, |\vec{k}| \sim q_0 \Rightarrow$  expand  $\sum_j T_j^{(h)}$  in  $D_h = \{k \in \mathbb{R}^d : |k_0| \gg |\vec{p}| \text{ or } |\vec{k}| \gg |\vec{p}|\}$
- **soft** ( $s$ ):  $k_0, |\vec{k}| \sim |\vec{p}| \Rightarrow$  expand  $\sum_j T_j^{(s)}$  in  $D_s = \{k \in \mathbb{R}^d : |\vec{k}| \lesssim |k_0| \lesssim |\vec{p}|\}$
- **potential** ( $p$ ):  $k_0 \sim \frac{\vec{p}^2}{q_0}, |\vec{k}| \sim |\vec{p}| \Rightarrow$  expand  $\sum_j T_j^{(p)}$  in  $D_p = \{k \in \mathbb{R}^d : |k_0| \ll |\vec{k}| \lesssim |\vec{p}|\}$

[no explicit boundaries needed]

$$\hookrightarrow D_h \cup D_s \cup D_p = \mathbb{R}^d, \quad D_h \cap D_s = D_h \cap D_p = D_s \cap D_p = \emptyset$$

$\hookrightarrow$  The expansions  $T^{(h)}, T^{(s)}, T^{(p)}$  commute with each other.